BEST AVAILABLE COPY

For official use

14 JUN 199

17JUN 91H00407330 PAT 1.77 UC

Your reference

PH. 36402/m/P.

9112859.5

15.00

Notes

Ple type, or write in dark ink using CAPITAL letters. A prescribed fee is payable for a request for grant of a patent. For details, please contact the Patent Office (telephone 071–829 6910).

Rule 16 of the Patents Rules 1990 is the main rule governing the completion and filing of this form.

Do not give trading styles, for example, 'Trading as XYZ company', nationality or former names, for example, 'formerly (known as) ABC Ltd' as these are not required.

Patent Office

Request for grant of a Patent

Form 1/77

Patents Act 1977

Title of invention

1 Please give the title of the invention

PEPTIDE PROCESS

Applicant's details

- ☐ First or only applicant
- 2a If you are applying as a corporate body please give:

 Corporate name IMPERIAL CHEMICAL INDUSTRIES PLC

Country (and State United Kingdom of incorporation, if appropriate)

Warning

After an application for a Patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977 and will inform the applicant if such prohibition or restriction is necessary. Applicants resident in the United Kingdom are also reminded that under Section 23, applications may not be filed abroad without written permission unless an application has been filed not less than 6 weeks previously in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction revoked.

.

2b If you are applying as an individual or one of a partnership please give in full:

Surname

Forenames

2c In all cases, please give the following details:

Address Imperial Chemical House, Millbank,
London.

UK postcode

SW1P 3JF

(if applicable)

Country

United Kingdom

ADP number (if known)

935003

				•
	D 0: 1 :: .		٦	•
2d, 2e and 2f: If there are further	Second applicant 2d If you are applying	as a corporate body please give:	•	
applicants please provide details on a	Corporate name	and the service body process give.	1	
separate sheet of paper.	·			
	Courter to 18			
	Country (and State of incorporation, if			•
	appropriate)			
·			-	
	2e If you are applying	as an individual or one of a partnership please give in full:		
	Sumame			
	Forenames	•		
	2f in all cases pleas	e give the following details:	-	
		o ground removing double.		
	Address	•		
	UK postcode			
•	(if applicable)			
	Country			
	ADP number (if known)		·	
An address for service in the	Address for ser	vice details	-	
United Kingdom must be supplied		ed an agent to deal with your application?		
Please mark correct box	Yes 🔣 No	⇒ go to 3b		
	please give detail	is below	·	
•	Agent's name	REGINALD PETER SLATCHER		
,	Agent's address	Legal Department: Patents	į	
		Imperial Chemical Industries PLC		İ
		P O Box 6, Bessemer Road Welwyn Garden City		T. O. D. D. C. D.
		Hertfordshire		
	Postcode	AL7 1HD		
	Agent's ADP	01320837001 /	·	
	number	7		
3b: If you have appointed an agent, all	3b If you have not ap	pointed an agent please give a name and address in the		
correspondence concerning your application will be sent to the agent's	United Kingdom to	o which all correspondence will be sent:		
United Kingdom address.	Name			
	Address			
			Į.	
		Daytime telephone	1	!
	Postcode	number (if available)		
	ADP number (if known)			
			ī	

ſ	Reference number				
	Agent's or applicant's reference PH. 364-00/m/P				
	❸ Claiming an earlier application date				
	5 Are you claiming that this application be treated as having been filed on the date of filing of an earlier application?				
Please mark correct box	Yes No X → go to 6 please give details below				
•	number of earlier				
	application or patent number				
	iday month year)				
•					
	and the Section of the Patents Act 1977 under which you are claiming:				
Please mark correct box	15(4) (Divisional) 8(3) 12(6) 37(4)				
6 If you are declaring priority from a PCT Application please enter 'PCT' as	Declaration of priority				
the country and enter the country code (for example, GB) as part of the	6 If you are declaring priority from previous application(s), please give:				
	Priority application number Filing date				
application number.	Country of filing (if known) (day, month, year)				
Please give the date in all number format, for example, 31/05/90 for 31 May 1990.					
Please give the date in all number format, for example, 31/05/90 for					
Please give the date in all number format, for example, 31/05/90 for					
Please give the date in all number format, for example, 31/05/90 for					
Please give the date in all number format, for example, 31/05/90 for					
Please give the date in all number format, for example, 31/05/90 for					
Please give the date in all number format, for example, 31/05/90 for					
Please give the date in all number format, for example, 31/05/90 for					
Please give the date in all number format, for example, 31/05/90 for					
Please give the date in all number format, for example, 31/05/90 for					
Please give the date in all number format, for example, 31/05/90 for					

The answer must be 'No' if: any applicant is not an inventor there is an inventor who is not an	InventorshipAre you (the applicant or applicants) the sole inventor or the joint inventors?
applicant, or any applicant is a corporate body.	Please mark correct box Yes No X Porm 7/77 will need to be filed (see Rule 15).
Please supply duplicates of claim(s), abstract, description and	© Checklist
drawing(s).	8a Please fill in the number of sheets for each of the following types of document contained in this application.
•	Continuation sheets for this Patents Form 1/77
	Claim(s) Description 6
	Abstract Drawing(s)
	8b Which of the following documents also accompanies the application?
:	Priority documents (please state how many)
:	Translation(s) of Priority documents (please state how many)
; -	Patents Form 7/77 - Statement of Inventorship and Right to Grant (please state how many)
Please mark correct box(es)	Patents Form 9/77 – Preliminary Examination/Search
	Patents Form 10/77 – Request for Substantive Examination
You or your appointed agent (see Rule 90 of the Patents Rules 1990)	Request
must sign this request.	I/We request the grant of a patent on the basis of this application. IMPERIAL CHEMICAL INDUSTRIES PLC
Please sign here 📦	Signed of Date 1/4 /6/91.
	AUTHORISED OFFICER
A completed fee sheet should preferably accompany the fee.	Please return the completed form, attachments and duplicates where requested, together with the prescribed fee to:
: : :	☐ The Comptroller The Patent Office State House 66–71 High Holborn
	London WC1R 4TP

- 1 -

PEPTIDE PROCESS

This invention relates to a process for making peptides and more particularly it relates to a solid phase peptide synthesis method for the preparation, inter alia, of the decapeptide goserelin.

The solid phase synthesis of peptides has been known for almost 30 years following the pioneering work of Merrifield first published in 1962. The general principle of this type of synthesis is as follows:-

- (a) An N-protected amino acid (the protecting group is commonly t-butoxycarbonyl, abbreviated to Boc) is attached to a solid, non-soluble support (commonly a polystyrene resin) at its carboxylic end via a linking group (commonly a benzyl ester).
- (b) The \underline{N} -protecting group is removed by means which do not detatch the amino acid from the solid support, and a second \underline{N} -protected amino acid is coupled to the one already attached (commonly by use of a carbodi-imide coupling agent).
- (c) The sequence is repeated using as many N-protected amino acids as are required until the desired peptide has been formed, still attached at its carboxyl end to the solid support.
- (d) The final N-protecting group is removed and the peptide is separated from the solid support by cleavage of the linking group (commonly by use of a strong acid).

The whole synthesis can be machine-aided and in some circumstances the peptide may be formed without manual intervention. The Boc protecting groups are removed by triflouroacetic acid and the peptide chain is removed from the solid support with a stronger acid such as hydrofluoric acid.

Since the introduction of this technique many modifications have been introduced, but the process is essentially as first proposed. Two major

innovations have been the use of a polyamide as the solid support and the use of a N-fluoren-9-ylmethoxycarbonyl (Fmoc) protecting group for the N^{α} -group of the amino acid. The Fmoc group is distinguished by being labile to base (commonly piperidine). For further detail reference is made, for example, to Atherton and Sheppard, "Solid phase peptide synthesis - a practical approach", IRL Press at Oxford University Press, 1989; Barany et al., "Solid-phase peptide synthesis: a silver anniversary report", Int. J. Peptide Protein Res., 1987, 30, 705-739 and Fields et al., ibid, 1990, 35, 161-214.

Throughout this specification standard abbreviations for amino acids, protecting groups, coupling agents and the like will be used. For the avoidance of doubt, as well as Boc and Fmoc defined above, the following are relevant standard abbreviations:-

Arg arginine

Azgly azaglycine (H2N-NH-C00H)

D-Ser D-serine

Glp pyroglutamic acid

His histidine

Leu leucine

Pro proline

Ser serine

Trp tryptophan

Tyr tyrosine

DIPC di-isopropylcarbodi-imide

HOBt 1-hydroxybenzotriazole

DMF N.N-dimethylformamide

BrZ 2-bromobenzyloxycarbonyl

Bu^t tert-butyl

Bzl benzyl

Goserelin is an LHRH analogue used in the treatment of prostate cancer, breastcancer and certain gynaecological conditions. In the first-mentioned treatment it acts by inducing a chemical castration.

Its structure is:-

${\tt Glp-His-Trp-Ser-Tyr-D-Ser(Bu}^{\sf t})-{\tt Leu-Arg-Pro-Azgly-NH}_2$

It will be seen that there are two features of this structure which are incompatible with traditional solid phase peptide synthetic routes. The first is the Azgly carboxy terminal amino acid; procedures for linking such a group to a solid support are not known. Free azaglycine has a terminal -NH-COOH group, which is an unstable carbamic acid.

The second is the t-butyl group attached to the D-serine moeity; in order to preserve this group traditional means for removing the completed peptide from the solid support cannot be used.

We have now found a method of preparing goserelin and similar peptides by solid phase synthesis.

According to the invention there is provided a method for solid phase synthesis of a peptide containing a C-terminal aza-amino acid amide, which comprises

- (i) assembling all the amino acids of the peptide except the C-terminal aza-amino acid by conventional solid phase synthesis;
- (ii) cleaving the peptide from the support with hydrazine or a substituted hydrazine; and
- (iii) reacting the hydrazide thus released with a cyanate ion.

The last two stages of this process form firstly a peptide with the carboxyl end of the formula:-

-CONH-NH-R

wherein R is hydrogen (in azgly) or such a group that H₂N-NR-COOH is an aza-analogue of an amino acid, and secondly a peptide with the carboxyl end of the formula:-

-CONH-NR-CONH,

The cleavage of the peptide from the support may be carried out using hydrazine or a substituted hydrazine in solution in DMF, N-methylpyrrolidone or a similar solvent.

A suitable cyanate ion may be provided by an alkali metal cyanate, for example potassium cyanate. The reaction may be carried out in aqueous acidic conditions.

According to a further feature of the invention there is provided a method for solid phase synthesis of a peptide containing an amino acid which contains a t-butyloxy group in its sidechain which comprises the use of a linking group connecting the amino acid to the solid support which is labile under conditions which do not cleave an <u>0</u>-t-butyl group.

A suitable linking group is one which may be cleaved by the use of hydrazine which will not cleave the t-butyl ether.

The amino acids contained in such a peptide are the t-butyl ethers of, for example, serine, D-serine, threonine, tyrosine and hydroxyproline.

The invention is illustrated but not limited by the following example:-

Example

(a) Solid phase perparation of nonapeptide

The solid phase synthesis was carried out in automatic mode on an Applied Biosystems 430A Peptide Synthesizer using Boc-Pro-OBzl-polystyrene resin 1% cross-linked with divinylbenzene (Peninsula Laboratories, 1.25g, 0.38 meq/g though nominally 0.7 meq/g). The following protected amino acids were converted to benzotriazolyl esters by reaction with HOBt and DIPC in DMF immediately before use. The protected amino acids were coupled in the following sequence:-

Boc-Arg(HCl)-OH
Boc-Leu-OH
Fmoc-DSer(Bu^t)-OH
Fmoc-Tyr(BrZ)-OH
Fmoc-Ser-OH
Fmoc-Trp-OH
Fmoc-His(Fmoc)-OH
Pyr-OH

The sequence of operations for the first two stages (using Boc-protected- amino acids) was:-

removal of Boc with 45% triflouroacetic acid in methylene chloride 10% DIEA/DMF wash coupling (2 equivalents of protected amino acid HOBt ester) removal of Boc as above

The sequence of operations for the last six stages (using Fmoc-protected-amino acids) was:-

removal of Fmoc with 20% piperidine/DMF 0.5 molar HOBt/DMF wash coupling (1 equivalent of protected amino acid HOBt ester)

All coupling reactions except that using Boc-Arg(HCl)-OH were of 1 hour duration; the Boc-Arg(HCl)-OH one was of 2 hours duration. There was thus obtained the nonapeptide-resin (1.7g; 0.29 mmole peptide per g.) with the Tyr still protected by BrZ.

(b) Cleavage of peptide from resin

The peptide resin prepared above was treated with a 20-fold excess of anhydrous hydrazine in DMF (20ml) at laboratory temperature for 24 hours, and the mixture was filtered and evaporated to dryness. This procedure also removed the BrZ protecting group from the Tyr moeity.

The residue was purified by gel filtration on a column (LH 20 Sephadex) using a 20:1 v/v mixture of water and acetic acid as eluant. There was thus obtained

 ${\tt Pyr-His-Trp-Ser-Tyr-DSer(Bu}^t)-{\tt Leu-Arg(H}^+)-{\tt Pro-NH-NH}_2$

the structure of which was confirmed by amino acid analysis and FAB mass spectroscopy $(M+H)^{+} = 1226$.

(c) Preparation of goserelin

A solution of potassium cyanate (11mg) in water (1.36ml) was added portionwise during 1 hour to a solution of the above hydrazide (118mg) in a 20:1 v/v mixture of water and acetic acid (10ml). The mixture was freeze-dried and the residue was purified by reverse-phase column chromatography (Dynamax 60Å, C₁₈, 1 inch diameter) using a gradient of 10% to 40% by volume of acetonitrile in water containing 0.1% trifluoroacetic acid. There was thus obtained goserelin (100mg, 25% yield overall), the structure of which was confirmed by FAB mass spectroscopy.

PP 91/58 13JUN91



This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

D BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
☐ OTHER:

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.